

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/86, C07K 14/535	A2	(11) International Publication Number: WO 96/09399 (43) International Publication Date: 28 March 1996 (28.03.96)
(21) International Application Number: PCT/US95/11537 (22) International Filing Date: 12 September 1995 (12.09.95) (30) Priority Data: 311,485 23 September 1994 (23.09.94) US (71) Applicant: SOMATIX THERAPY CORPORATION [US/US]; Suite 100, 950 Marina Village Parkway, Alameda, CA 94501 (US). (72) Inventors: SHANKARA, Srinivas; Apartment E, 2255 San Jose Avenue, Alameda, CA 94501 (US). DWARKI, Varavani; Apartment N, 1175 Broadway Street, Alameda, CA 94501 (US). NIJJAR, Tarlochan; 946 Foxfire Drive, Manteca, CA 95336 (US). (74) Agents: HALLUIN, Albert, P. et al.; Pennie & Edmonds, 1155 Avenue of the Americas, New York, NY 10036 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>

(54) Title: CHIMERIC ADENOVIRUS FOR GENE DELIVERY**(57) Abstract**

Chimeric adenovirus capable of transducing mammalian cells with DNA of interest are disclosed. The chimeric adenovirus are useful for the delivery of cloned genes into an individual and are therefore also useful for treating mammalian genetic diseases and disorders.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

CHIMERIC ADENOVIRUS FOR GENE DELIVERY

1. FIELD OF THE INVENTION

5 The present invention is directed to novel adenovirus vectors useful for the delivery of cloned genetic material to target cells. The chimeric adenovirus vectors comprise genetic material of interest which is flanked by adenoviral sequences, and may optionally comprise a suitable eucaryotic
10 promoter to facilitate the expression of the genetic material of interest. The chimeric adenovirus are produced by a process involving a recombinant adenovirus vector which is used in conjunction with replication deficient helper adenovirus genomes to generate recombinantly produced chimeric
15 adenovirus particles comprising the genetic material of interest. The resulting chimeric adenovirus may be used to infect target cells which subsequently express the cloned genetic material. One class of novel chimeric adenovirus does not contain a selectable marker which obviates the need for a
20 selection step after the genetic material of interest has been introduced into the target cells.

2. BACKGROUND OF THE INVENTION

Mammalian cells may be transduced by any of a variety of
25 well known processes. Techniques such as calcium phosphate precipitation and DEAE-dextran mediated transfection are widely used in the art. More recently, other techniques for delivery of exogenous DNA into cells such as electroporation or the use of liposomes have gained increased acceptance.
30 Perhaps the most elegant methods of introducing recombinant nucleic acid into cells is viral mediated cell transduction.

Recombinant retroviruses have been widely used in gene transfer experiments (see generally, Mulligan, R.C., Chapter 8, In: Experimental Manipulation of Gene Expression, Academic
35 Press, pp. 155-173 (1983); Coffin, J., In: RNA Tumor Viruses, Weiss, R. et al. (eds.), Cold Spring Harbor Laboratory, Vol. 2, pp. 36-38 (1985). Other eucaryotic viruses which have been

used as vectors to transduce mammalian cells include adenovirus, papilloma virus, herpes virus, adeno-associated virus, rabies virus, and the like (See generally, Sambrook et al., Molecular Cloning, Cold Spring Harbor Laboratory Press, 5 Cold Spring Harbor, New York, Vol. 3:16.1-16.89 (1989).

Adenovirus have proved to be of particular interest because of several features of adenoviral biology (See generally, Berkner, K.L. (1992) Curr. Top. Microbiol. Immunol. 158:39-66). For instance, viral concentration, or titer, may
10 be an important factor in achieving high efficiency transduction of mammalian cells. Adenovirus, by virtue of their life-style, generally allow growth conditions which result in production of higher titer stocks than other mammalian virus.

15 Also unlike other viruses, adenovirus capsids are not enveloped. Because of this fact, adenovirus particles are quite stable, and may retain infectivity after any of a variety of laboratory procedures. Procedures of particular interest include methods of concentrating infective virus,
20 e.g., CsCl centrifugation, or methods that allow virus to be stored for relatively long periods while retaining substantial infectivity.

Furthermore, the expression of genes encoded by recombinant adenovirus does not require target cell
25 proliferation or viral integration, although a small subset of the adenovirus presumably integrate into the host genome during infection. Hence, adenoviral vectors are generally better suited than other viral vectors for the transduction of postmitotic, slowly proliferating, or nonreplicating cells.

30 Additionally, particularly where species-specific infection is preferred, replication deficient human, or murine, adenovirus are available for the construction of recombinant virus particles that express a gene of interest. Thus, unlike transduction systems using other eucaryotic virus
35 vectors, recombinant adenovirus can be engineered to utilize viral coat proteins which normally facilitate the normal infection of human cells or cells of other species, rather

then rely on the viral coats of a less specific, or amphotropic, nature. This species specificity appears to result in more efficient infection kinetics than can generally be obtained by virus with less specific infectivity.

5 An additional advantage of using adenovirus for gene delivery is that the genetic material transduced (to be expressed) into the host cell is DNA. Thus, expression of the transduced gene does not need to be preceded by reverse transcription. This is particularly advantageous where the
10 intended recipient is undergoing treatment for the suppression of retroviral disease (i.e., AZT treatment to inhibit reverse transcriptase activity), such as treatment for acquired immunodeficiency syndrome (AIDS).

Recombinant adenoviral vectors have been generated which
15 express a variety of genes. Perhaps most notable is the replication deficient adenovirus vector Ad.RSV that expresses incorporated genetic material of interest using an incorporated promoter from the Rous Sarcoma Virus. In particular, Ad.RSV beta gal (which expresses the bacterial β -
20 galactosidase gene) has been used as a marker for in vivo gene transfer experiments involving salivary glands (Mastrangeli et al. (1994) Am. J. Physiol. 266:1146-1155); mesothelial cells (Setoguchi et al. (1994) Am. J. Respir. Cell. Mol. Biol. 10(4):369-377); and tumor cells (Brody et al. (1994) Hum. Gene
25 Ther. 5(4):437-447, Chen et al. (1994) Proc. Natl. Acad. Sci., U.S.A. 91(8):3054-3057).

An ideal replication deficient adenovirus for the delivery of genetic material of interest would comprise a variety of structural and functional elements. It would
30 readily infect target cells of interest; it would place the gene of interest under the control of a well-characterized eucaryotic promoter element; it would create a gene structure flanking the gene of interest which would provide properly spaced and oriented genetic elements to allow optimum
35 translational efficiency and mRNA stability; and it would produce high titer and substantially helper-free stocks of the recombinant adenovirus.

3. SUMMARY OF THE INVENTION

The present invention relates to replication deficient chimeric adenovirus that allow for the rapid insertion and expression of deoxyribonucleic acid (DNA) of interest into mammalian cells, either in vitro or in vivo. The DNA of interest can optionally comprise a gene, or fraction thereof, oriented to express either a polypeptide or protein of interest, or a "sense" or "antisense" nucleic acid of structural or regulatory importance. Preferably, the DNA of interest will be placed in an expression cassette that contains a eucaryotic promoter and/or enhancer region; nucleotide sequence corresponding to a retroviral Psi-packaging site; and a substantially noncoding 3' DNA which facilitates the stability, polyadenylation, or splicing of the transcript.

The chimeric adenovirus are thus useful for both the transduction of mammalian cells, and the expression of DNA of interest to produce regulatory factors or proteins. The regulatory factors or proteins may optionally be produced in culture or otherwise such that they can be subsequently purified and used for therapeutic, medicinal or diagnostic purposes.

The chimeric adenovirus are particularly useful for gene therapy, replacement, or insertion because of the high infectivity inherent in adenovirus biology; the high viral concentrations which may be produced during the culture and subsequent concentration of the chimeric adenovirus; and the relatively long storage life of the chimeric particles.

Either murine, or human adenovirus of serotypes A, B, or C may be used in the present invention. Of particular interest are type C adenovirus (used in the present invention) which retain infectivity while generally being considered nononcogenic.

4. BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a schematic representation of the method of producing chimeric adenovirus via the recombination of

cotransfected plasmids. One plasmid, pXCJL-GMCSF, contains a "cassette" comprising the gene encoding the cytokine granulocyte/macrophage colony stimulating factor (GMCSF) situated such that it is transcribed, processed, and
5 translated under the regulatory control of flanking viral sequences. The second plasmid, pJM17, comprises a replication and packaging deficient adenovirus "helper" genome. The two plasmids must recombine to produce a packagable genome, and thus substantially all of the resulting virus comprise the
10 chimeric adenovirus desired (Recombinant E1-deleted GM-CSF adenovirus).

Figure 2 presents a schematic diagram and partial restriction map of pJM17.

15

Figures 3A-E disclose the DNA sequence of pXJCL-hGM-CSF (SEQ. I.D. NO. 1), the plasmid used to construct the human GM-CSF expression cassette, and in the recombinatory insertion of the GM-CSF expression cassette into the replication deficient
20 genome contained in pJM17. The sequence of the murine GM-CSF is disclosed in foreign patent EP177568B1, herein incorporated by reference.

Figures 4A and 4B show the transient expression of human
25 GM-CSF after one month old Balb/c mice were intramuscularly injected with either 10^9 or 10^8 pfu of Ad.hGM-CSF respectively. Serum samples were taken up to twenty one days after infection and GM-CSF levels were assayed by ELISA. Individual mice are represented by number and correspond to
30 the indicated bars on the graphs.

Figure 5 shows the expression of human GM-CSF (as quantified by ELISA) after Ad.hGM-CSF injection and reinjection into adult Balb/C mice. Four month old Balb/C
35 mice were injected with 10^8 pfu of Ad.hGM-CSF either I.V. (mice 103 and 105) or I.M. (mice 201, 203, and 205). All mice were reinjected (I.M.) with 10^9 pfu of Ad.hGM-CSF at day 31.

Figure 6 shows the expression of human GM-CSF (as quantified by ELISA) after Ad.hGM-CSF injection and reinjection into adult SCID mice. SCID mice were injected (I.V.) with 10^8 pfu of Ad.hGM-CSF, and GM-CSF blood serum levels were subsequently monitored. All mice were reinjected (I.M.) with 10^8 pfu of Ad.hGM-CSF at day 31, and monitored for GM-CSF expression through day 71.

5. DETAILED DESCRIPTION OF THE INVENTION

- 10 The present invention provides for chimeric adenovirus which are useful for transducing mammalian cells with DNA of interest, as well as methods of producing and using the chimeric adenovirus. Previous recombinant adenovirus expression vectors have specifically taught the expression of
- 15 the genetic material of interest under the control of endogenous adenoviral promoters, or have suggested that the DNA of interest be inserted into recombinant adenovirus under the control of an RSV promoter already present in the vector Ad.RSV.
- 20 In the present system, the particular DNA of interest is first constructed as an expression cassette which comprises a gene, or portion thereof, of interest that is flanked by sequences of viral origin which are spatially organized to optimize the expression of the DNA of interest. As used
- 25 herein, the term "expression" refers to the transcription of the DNA of interest, and the splicing, processing, stability, and, optionally, translation of the corresponding mRNA transcript. The recombinant DNA cassette is subsequently recombined into a replication deficient helper adenovirus to
- 30 produce the infective chimeric adenovirus of interest. This method best ensures the maximal expression of the DNA of interest and additionally provides a method that is generally applicable to the relatively facile production of chimeric adenovirus which express a wide variety of DNAs.
- 35 The particular advantage of using an expression cassette stems from the fact that the recombinant Ad.RSV vector is rather large (over 36kb). This large size makes plasmids

which contain the Ad.RSV genome somewhat difficult to engineer as the number of unique (and hence useful) restriction sites tends to diminish as the amount of DNA sequence increases. Thus, the utilization of a smaller plasmid to construct the
5 expression cassette better enables a wide variety of genetic engineering techniques which may allow the fine tuning of the expression of the DNA of interest (see generally, Sambrook et al. (1989) Molecular Cloning Vols. I-III, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, and Current
10 Protocols in Molecular Biology (1989) John Wiley & Sons, all Vols. and periodic updates thereof, herein incorporated by reference). For instance, after the DNA of interest is placed between the desired regulatory elements (i.e., promoter and poly-adenylation signal), unwanted regions of extraneous DNA
15 may be looped-out and deleted by site-directed mutagenesis (Krogstad and Champoux (1990) J. Virol. 64(6):2796-2801, herein incorporated by reference) such that the DNA of interest is precisely placed relative to the promoter and splicing elements, and, if a protein or polypeptide is
20 desired, a strong Kozak translation start site (Kozak (1989) J. Cell Biol. 108:229). This arrangement best ensures that the resulting chimeric adenovirus will maximally express the DNA of interest.

As used herein, the term replication defective
25 adenovirus, refers to a adenovirus that are incapable of self replication within host cells that, absent infection or transfection, do not express at least one adenovirus gene or gene product.

Any number of transcriptional promoters and enhancers may
30 be used in the expression cassette, including, but not limited to, the herpes simplex thymidine kinase promoter, cytomegalovirus promoter/enhancer, SV40 promoters, and retroviral long terminal repeat (LTR) promoter/enhancers. Of special interest are any of a number of well characterized
35 retroviral promoters, particularly the Moloney murine leukemia virus (MLV) LTR promoter and the human immunodeficiency virus (HIV) LTR.

According to one embodiment of the present invention, recombinant DNA techniques have been used to construct expression cassettes in plasmid pXCJ1.1 which comprise genes coding for the murine or human forms of granulocyte macrophage colony stimulating factor (GM-CSF), which have been placed under the transcriptional control of the Moloney murine leukemia virus (MLV) long terminal repeat (LTR). In a further embodiment, an SV40 poly-adenylation sequence flanks the 3' end of the GM-CSF gene. Thus, the transcript produced by either GM-CSF expression cassette is transcribed using the MLV LTR promoter and enhancer sequences, poly-adenylated using an SV40 poly-adenylation sequence, spliced using the MLV splice donor and splice acceptor sequences, and the mRNA is presumably translated using the endogenous MLV translation initiation sequence of the MLV gag gene. By engineering the DNA expression cassette such that the resulting transcript surrounds the coding region with naturally occurring viral control sequences, near optimum mRNA stability is obtained. Thus, as used herein, the terms "DNA expression cassette" or simply "expression cassette" both refer to a DNA molecule comprising a eucaryotic promoter and/or enhancer region, a DNA of interest to be transcribed by the promoter, and a substantially noncoding 3' region of DNA that facilitates the stability, polyadenylation, or splicing of the transcript.

The GM-CSF expression cassette is inserted into a replication defective helper adenovirus via homologous recombination after two circular plasmids (one containing the GM-CSF expression cassette and the other containing the replication defective adenovirus genome) are co-transfected into the appropriate cell line (see Fig. 1). Using this system, only the specifically desired chimeric adenovirus are packaged. The resulting chimeric adenovirus expresses a mammalian gene (human or murine GM-CSF) that is expressed under the transcriptional and translational control of MLV and SV40 control sequences. The chimeric adenovirus can subsequently be purified by any of a number of well established techniques including, but not limited to, plaque

purification, purification by limiting dilution, or the like. Purified chimeric adenovirus can then be propagated to relatively high titers by infection of appropriate host cells, for example 293 cells (human kidney epithelial cells which constitutively produce adenovirus E1A). Although the chimeric adenovirus infections will generally produce highly concentrated viral preparations, one may elect to further concentrate and purify the chimeric adenovirus to achieve titers of about $1-5 \times 10^{11}$ plaque forming units (pfu)/ml by CsCl density equilibrium centrifugation (followed by dialysis), ultrafiltration, or the like.

The resulting chimeric adenovirus, designated Ad.mGM-CSF (murine GM-CSF) or Ad.hGM-CSF (human GM-CSF), have been shown to be useful for the production of microgram quantities (as quantified by enzyme linked immunosorbent assay, or ELISA) of GM-CSF in infected NIH 3T3 cells (see Table 1). The properties of Ad.hGM-CSF and Ad.mGM-CSF make both ideally suited for applications where GM-CSF expression by any of a broad range of target cells may be desired.

Of particular interest is the use of Ad.hGM-CSF or Ad.mGM-CSF to transduce primary tumor cells. It has previously been established that vaccinations with tumor cells engineered to secrete GM-CSF can stimulate anti-tumor immunity in mice (Dranoff et al. (1993) Proc. Natl. Acad. Sci. U.S.A. 90:3539-3543. Ad.hGM-CSF has been used to transduce primary human melanoma, renal cell carcinoma, and colon carcinoma cells which subsequently produced microgram quantities (about $1-5 \mu\text{g}/10^6$ cells) of human GM-CSF (see Tables 2a-d). Additionally, Ad.mGM-CSF has been used to infect and transduce murine B16 melanoma cells which may subsequently be irradiated (using about 5,000 rads) and assessed for efficacy as an anti-melanoma vaccine.

Ad.hGM-CSF was also injected into Balb/c or SCID mice at various anatomical locations, and in vivo expression of GM-CSF was detected and quantified by ELISA (see Figs. 5 & 6).

Ad.hGM-CSF has been deposited (received at the ATCC on September 23, 1994) at the American Type Culture Collection,

Rockville, MD, under the accession number _____ under the terms of the Budapest Treaty. Applicants further agree to make this deposit available, without restriction to responsible third parties upon the granting of a patent from 5 this application in the United States, and comply with existing laws and regulations pertaining thereto, without limitation, except as to third parties adherence to applicants rights as prescribed by the claims of a patent issuing from this application.

10

As described briefly above and in detail in the Examples, the present invention provides a method of producing chimeric adenovirus comprising the recombinatory insertion of a DNA expression cassette contained in a circular plasmid into a 15 replication deficient helper adenovirus genome contained in a circular plasmid to produce a chimeric adenovirus capable of transducing mammalian cells. The use of two circular plasmids is an important feature of the method of the present invention, since there is no need to linearize the adenoviral 20 helper genome prior to cotransfection.

The chimeric adenovirus of the present invention exhibit very high infectivity and thus high levels of cellular transduction and expression of a DNA of interest. In addition to the specifically disclosed GM-CSF genes, modified forms of 25 the GM-CSF genes may be utilized which have been altered by deletion or insertion, or to optimize codon usage for the specific target cells intended. DNA expression cassettes may also be constructed which allow the subsequent production of chimeric adenovirus which are capable of transducing any of a 30 number of heterologous mammalian genes (i.e., DNAs of interest, subject to the restriction that the net size of the insert is less the about 9 kb in length).

Besides GM-CSF, other heterologous genes of particular interest include, but are not limited to, nerve growth factor 35 (NGF), tyrosine hydroxylase (TH), ciliary neurotropic factor (CNTF), brain-derived neurotropic factor (BDNF), factors VIII and IX, tissue plasminogen activator (tPA), interleukins 1-2

and 4-6, tumor necrosis factor- α (TNF- α), α or γ interferons, and erythropoietin. Chimeric adenovirus that express any of the above genes, or portions thereof, may be particularly useful for the treatment of mammalian diseases or disorders
5 related to aberrant or deficient levels of the corresponding polypeptides or proteins in a given individual.

Alternatively, chimeric adenovirus containing the genes for these factors may also be used to generate transient expression of the factors in vivo as required to
10 therapeutically treat medical crisis. For instance, an infusion of chimeric adenovirus containing a tPA expression cassette would provide transient expression of tPA during the critical period following a heart-attack or stroke.

The high efficiency transduction inherent in the chimeric
15 adenovirus system makes them particularly well suited for the treatment of genetic or inherited disease, as well as the treatment of acquired disease. For instance, chimeric adenovirus may be used to deliver genes into a variety of cell types to correct genetic defects associated with diseases
20 including but not limited to β -thalassemia, phenylketonuria, sickle-cell anemia, cystic fibrosis, or adenosine deaminase deficiency.

The chimeric adenovirus of the present invention may be used to transduce mammalian cells either in vitro or in vivo.
25 Where transduction in vitro is contemplated, cells may be infected at multiplicities of infection (moi's) of between about 1:1 to about 5000:1, and generally in the range of about 100:1 to about 2,500:1. Moi's of up to about 1000:1 have produced good expression of the DNA of interest without
30 evidence of serious cellular toxicity effects, and moi's of about 200:1 have resulted in no toxicity. Using similar methodologies, chimeric adenovirus may be used to infect resected primary tissue or cells which may subsequently be reintroduced into the body of an individual by established
35 surgical or medical procedures.

Where diagnostic, therapeutic or medicinal use of chimeric adenovirus is contemplated, chimeric adenovirus

capable of transducing and expressing the DNA of interest may be introduced in vivo by any of a number of established methods. For instance, chimeric adenovirus may be administered by inhalation. Alternatively, chimeric
5 adenovirus suspensions may also administered by intravenous (I.V.), intraperitoneal (I.P.), or intramuscular (I.M.) injection.

The chimeric adenovirus may also be injected directly into tumors. To prove the feasibility of this concept, a
10 chimeric adenovirus which encodes a bacterial lacZ gene was injected into B16 melanoma tumors in C57 mice. Following injection, adenovirus mediated transduction and in vivo expression of β -galactosidase was observed in the tumors.

Other in vivo studies have established that a single
15 bolus of as much as about 10^9 pfu (in 100 μ l total volume) of Ad.hGM-CSF can be injected (I.V. or I.M.) into mice without apparent toxicity effects (see Fig. 4A).

Possible cell types or tissues that may serve as targets for chimeric adenovirus gene delivery include, but are not
20 limited to, hepatocytes, fibroblasts, endothelial cells, bone marrow stem cells, lymphocytes, neural tissue, astrocytes, alveolar tissue, and granulocytes.

An additional embodiment of the present invention is chimeric adenovirus containing expression cassettes which
25 further comprise a specific retroviral Psi-packaging sequence. More particularly, a Psi-packaging sequence which corresponds to that recognized and used by any of a number of ecotropic and amphotropic Moloney murine leukemia virus packaging cell lines including, but not limited to, PA317 or PsiCRIP.

30 Where the above expression cassette of the chimeric adenovirus further encodes at least a portion of an MLV 3' LTR sequence (minimally comprising the U3 and R regions of the LTR) located distal to the gene of interest, the chimeric adenovirus may be used to transiently infect MLV packaging
35 cell lines and produce amphotropic or ecotropic retrovirus which package RNA genomes transcribed by the expression cassette of the chimeric adenovirus. Infection of the

appropriate cells by the resulting retrovirally packaged chimeric adenovirus transcripts will result in the integration and stable expression of the DNA of interest contained in the expression cassette of the chimeric adenovirus. The chimeric
5 adenovirus described above provide the user with increased versatility relative to previously disclosed retroviral or adenoviral transduction vectors. This is because a single chimeric adenovirus allows the user to choose between the increased storage life, infectivity, and transient expression
10 inherent in the high titer chimeric adenovirus system, or the stable integration and expression inherent in the MLV packaging system. Alternatively, an optimal mixture of the two delivery systems may be preferred. Thus, the present invention also provides for replication defective chimeric
15 adenovirus which contain an expression cassette which further comprises nucleotide sequence corresponding to a MLV Psi-packaging site.

An additional embodiment of the present invention is chimeric adenovirus which place the expression of genes whose
20 products are toxic to the cell under the strict control of a trans-activated promoter, such as an HIV LTR promoter. Toxic genes which may be employed in these vectors include, but are not limited to, sequence coding for diphtheria toxin A chain, polio virus protein 2A, and the like (or modified forms
25 thereof). Since the HIV promoter generally requires virally encoded trans- activators, chimeric adenovirus will generally only express the toxic products (hence killing the cells) in HIV infected cells. Thus, since the expression of genes contained in chimeric adenovirus is not dependent on cell
30 division or proliferation (unlike retrovirally expressed genes), the above chimeric adenovirus may find utility in targeting and killing non-replicating or quiescent HIV-infected cells.

The present invention will now be illustrated by the
35 following examples, which are not intended to be limiting in any way.

6. EXAMPLES

6.1. CONSTRUCTION OF THE PXCJL-GMCSF PLASMID

The starting plasmid, designated PXCJL1, was constructed
5 from a modified Ad5 adenovirus genome cloned into pBR322. A
deletion was made from the map units 1.3 to 9.3, and a
multiple cloning site was inserted at the unique XbaI site.
This construct was obtained from Dr. Frank Graham of McMaster
University (McGrory, W.J. et al., Virology 163: 614-617,
10 1988).

The cDNA for human GM-CSF, along with upstream packaging
and splicing sequences and the complete MLV 5' LTR, were
isolated from plasmid MFGs-GM-CSF. MFGs is an unpublished
three nucleotide modification of the MFG vector, as
15 represented by MFG-GM-CSF (Dranoff, et al., Proc. Natl. Acad.
Sci. 90:3539-3543, 1993; the modification has no effect on
expression levels or transduction efficiencies). MFGs-GM-CSF
DNA was first digested to completion with HindIII and BamHI
and the ends were blunt-ended with the Klenow fragment. The
20 plasmid fragments were separated by electrophoresis on a 1%
agarose gel, and the 2.7 kb fragment extending from the 5' LTR
to the 3' end of the GM-CSF cDNA was purified from the gel
(Fragment 1).

The GM-CSF cDNA and associated sequences were then
25 subcloned into the multiple cloning site of PXCJL1 using
standard techniques (Sambrook, et al. Molecular Cloning: A
Laboratory Manual (1989)). The PXCJL1 plasmid was digested to
completion with XbaI, the ends were blunt-ended (end-filled)
with Klenow and treated with bacterial alkaline phosphatase.
30 This linearized vector fragment was purified from a 1% agarose
gel following electrophoresis (Fragment 2). The purified GM-
CSF cDNA (Fragment 1) was blunt-end ligated to the linearized
PXCJL1 with T4 ligase to generate the intermediate plasmid
PXCJL GM-CSF(I). XbaI and BamHI sites were regenerated in the
35 intermediate plasmid only if the insert was in the correct
orientation, as determined by restriction endonuclease (EcoRI
and BamHI) analysis.

To insert the SV40 polyadenylation sequence at the 3' end of the GM-CSF cDNA, PXCJL GM-CSF(I) was digested with BamHI and SalI, and the linearized fragment was isolated from a 1% agarose gel following electrophoresis (Fragment 3). The SV40 polyadenylation sequence was generated by polymerase chain reaction (PCR) using the pRC/CMV vector as the DNA template. The PCR primers were designed as follows:

the sense primer containing the BamHI site-
GAG GAT CCT ATC GCC TTC TTG ACG
10 and the antisense primer containing the SalI site-
GAG TCG ACT AAA CAA GTT GGG GTG.

PCR conditions were 95°C for 1 min., 55°C for 2 min., and 72°C for 3 minutes, for 35 cycles. The PCR product was cloned into a TA plasmid and sequenced. The product with the correct SV40 poly(A) sequence was digested with BamHI and SalI and the 216 bp SV40 poly(A) sequence was ligated to PXCJL GM-CSF(I) (Fragment 3) with T4 ligase.

The resulting cDNA expression plasmid, PXCJL, GM-CSF, contains the entire GM-CSF cassette, including the 5' MLV LTR, Psi-packaging and splicing sequences, the GM-CSF cDNA, and the SV40 poly (A) sequences, flanked by adenovirus sequences. Both murine and human GM-CSF cDNA were subcloned into PXCJL1 following the same strategy.

25 6.2. TRANSFECTION AND ISOLATION OF RECOMBINANT VIRUS

To generate recombinant virus, a replication deficient form of the adenoviral genome in circular form (pJM17) was obtained from Dr. Frank Graham. Techniques for transfection of 293 cells (a human kidney epithelial cell line), overlaying plates with agar-containing medium, picking and analysis of recombinant virus clones were carried out following the methods described by Graham and Prevec ("Manipulation of Adenovirus Vectors", in Gene Transfer and Expression Protocols, E.J. Murray, ed.). Briefly, 293 cells in 100 mm dishes were co-transfected with 10µg of pJM17 and 15µg of PXCJL-GMCSF plasmid by the calcium phosphate method following the standard transfection protocol. 36 hours after

transfection, cells were overlaid with 0.8% Noble agar containing DMEM with 10% heat inactivated fetal calf serum.

Plaques visible by 8 days after transfection were picked and resuspended in 1 ml of medium and freeze-thawed three times to release the virus. These supernatants were used as viral lysates in subsequent experiments. 0.2 ml of the viral supernatant from each individual plaque was added to the 1 ml of medium and used to infect confluent monolayers of 293 cells in a 6-well plate for four hours. After 24 hours, the cells began to show complete cytopathic effects.

At this time the colonies were harvested, and the medium was analyzed for GM-CSF secretion. The cells were lysed by three rounds of freeze-thaw, and the medium was used to infect NIH 3T3 cells in a 6-well plate. 80% confluent monolayers of NIH 3T3 cells in a 6-well plate were infected with 0.1 ml of crude virus stock in 1 ml of medium for four hours. 24 hours after infection fresh growth medium was added, and the GM-CSF secreted for the next 24 hours was analyzed by ELISA. The values for GM-CSF produced by Ad/human GM-CSF and Ad/mouse GM-CSF-transduced NIH 3T3 cells ranged from 300-400ng in 24 hours.

A schematic diagram of the recombination protocol used to generate Ad.hGM-CSF and Ad.mGM-CSF is presented in Figure 1.

6.3. PLAQUE PURIFICATION OF RECOMBINANT VIRUS

Confluent monolayers of 293 cells in 100mm dishes plated on day 1 were infected in 5 ml of medium on day 2 with 0.1 ml of viral supernatant obtained by resuspending virus containing agar block, as described above. After 1 hour of infecting at 37°C, the virus-containing medium was removed and overlaid with the agar-containing medium that had been prepared earlier. The cells were incubated at 37°C for 4-5 days and well isolated plaques were picked and analyzed for the ability to transduce NIH 3T3 cells with GM-CSF, as described earlier.

6.4. PURIFICATION AND AMPLIFICATION OF CHIMERIC ADENOVIRUS

Concentrated virus stocks were prepared from infected 293 cells. Confluent monolayers of 293 cells in 150mm dishes were infected with 5-10pfu/cell and after 36 hours when all the cells began to exhibit complete CPE, the cells were collected and resuspended in 5 ml of 0.1M Tris, pH 8.0. The virus was released from the cell pellets by three freeze-thaw cycles. After sonicating the cell lysate, 1.8 ml of saturated cesium chloride (in 10mM Tris, pH 8.0, 1 mM EDTA) was added to 3.1 ml of the cell lysate. This was centrifuged at 30,000 rpm in a SW 41 rotor for 20 hours. The virus band was collected and repurified by CsCl banding. The purified virus was then dialyzed against 10mM Tris/1 mM MgCl₂, pH 7.4, and stored in 10% glycerol at -70°C.

15

6.5. TRANSDUCTION OF NIH 3T3 CELLS WITH Ad.hGM-CSF AND Ad.mGM-CSF

NIH 3T3 cells were infected with purified virus at different multiplicities of infection (moi) for four hours, supernatants from 24-48 hours post-infection were collected and GM-CSF secretion was measured by ELISA. Results are shown in Table 1.

20

Table 1. Expression of human GM-CSF ($\mu\text{g}/1 \times 10^6$ cells/24 hr) in 3T3 cells.

25

TABLE 1.

moi	500	250	100	50
Ad.hGM-CSF	2.1	1.4	0.41	0.125
Ad.mGM-CSF	1.6	0.9	0.375	0.08

30

6.6. TRANSDUCTION OF PRIMARY HUMAN TUMOR CELLS WITH Ad.hGM-CSF Virus

Primary cultures of human melanoma, renal cell carcinoma, colon carcinoma and colorectal tumor cells were established and were transduced with Ad.hGM-CSF virus. The cultures were infected with Ad.hGM-CSF at different moi's for 4-8 hours,

35

supernatants were collected at 24-48 hours post-infection, and GM-CSF secretion was measured by ELISA. Results for the various cell types are presented in Tables 2a-d.

- 5 Tables 2a-d. Expression of GM-CSF ($\mu\text{g}/1 \times 10^6$ cells/24 hour) in Ad.hGM-CSF transduced primary tumor cells.

TABLE 2a.

10

moi	5000	1000	500	250	125	62.5	50
Melanoma-1 (P2)	2.3	12.6	5.4				1.1
Melanoma-2 (P2)		9.4	3.2	1.8	0.93	0.47	
Melanoma-3 (P2)		2.4	2.4	0.09	0.09	0.045	

TABLE 2b.

15

moi	5000	2500	1000	500	100
Renal Cell carcinoma (P3)	4.1	6.7	7.5	4.7	2.1

TABLE 2c.

20

moi	1000	200	100	20	10
Colorectal cells (P1)	0.15	1.8	1.5	0.42	0.22

TABLE 2d.

25

moi	5000	1000	500	50
Colon carcinoma (P1)	13.8	23.6	6.7	0.9

30

By comparison, transduction of the same types of human tumor cells by recombinant retrovirus expressing human GM-CSF results in expression in the range of 40-500 ng/ 1×10^6 cells/24 hours.

35

6.7. DELIVERY OF HUMAN GM-CSF INTO BALB/C MICE

To test for the ability of Ad.hGM-CSF to transduce mammalian cells in vivo, one month old Balb/C mice were injected intramuscularly (thigh muscle) with 100 μ l of purified virus at a concentration of either 10¹⁰ or 10⁹ pfu/ml. Transient expression of human GM-CSF was quantified by ELISA of serum samples taken from the mice at 2, 5, 7, 9, 14, and 21 days post infection. The data are presented in Figures 4A and 4B. Mice injected with 10⁹ pfu (Fig. 4A) exhibited peak expression of human GM-CSF five days after injection with transient expression tapering down to undetectable levels between seven to nine days after injection. Mice injected with 10⁸ pfu (Fig. 4B) also showed peak expression at about five days post injection but continued to express human GM-CSF until between nine to fourteen days after injection. These data clearly indicate that Ad.hGM-CSF transduces cells in vivo, and further mediates transient expression of human GM-CSF.

6.8. REPEATED INJECTION OF Ad.hGM-CSF INTO ADULT BALB/C MICE

To test whether Ad.hGM-CSF could also mediate transient expression of human GM-CSF in adult mice, and whether or not the route of injection substantially affected expression, four month old Balb/C mice were injected with 10⁸ pfu of Ad.hGM-CSF either intravenously (I.V.) or intramuscularly (I.M.). Serum samples were drawn at 3, 7, 14, and 31 days after injection and assayed for GM-CSF levels by ELISA. Serum levels of GM-CSF were generally lower than those observed in one month old mice, peaked between three to seven days after injection, and were undetectable fourteen days after infection.

Thirty one days after the initial injection the mice were reinjected (I.M.) with 10⁹ pfu of Ad.hGM-CSF and serum samples were drawn and analyzed for GM-CSF at 2, 4, and 9 days after reinjection. After reinjection, serum levels of GM-CSF peaked after two days and were undetectable after four days. The mode of primary injection apparently made little difference (see Fig. 5).

6.9. REPEATED INJECTION OF Ad.hGM-CSF INTO SCID MICE

To test whether an immune response might be the cause of the reduced expression of GM-CSF after reinjection, experiment 6.8 was essentially repeated using SCID (severe combined immunodeficiency) mice with the exception that Ad.hGM-CSF were only administered I.V.. As can be seen in Figure 6, SCID mice continued to express GM-CSF up to twenty eight days after initial infection and forty three days after I.M. reinjection of 10^9 pfu of Ad.hGM-CSF. These data (presented in Fig. 6) indicate that the diminution of GM-CSF levels in adult Balb/C mice seen in experiment 6.8 may be due to immune reaction to the adenovirus antigens expressed by the replication deficient genome of Ad.hGM-CSF.

All publications and patents mentioned in the above specification are herein incorporated by reference. The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the virus deposited since the deposited embodiment is intended as a simple illustration of one aspect of the invention and any virus that are functionally equivalent are within the scope of this invention. Various modifications of the invention in addition to those specifically shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

30

35

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Srinivas, Shankara
Dwarki, Varavani
Nijjar, Tarlochan
- (ii) TITLE OF INVENTION: Chimeric Adenovirus for Gene Delivery
- (iii) NUMBER OF SEQUENCES: 1
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Pennie & Edmonds
 - (B) STREET: 2730 Sand Hill Road
 - (C) CITY: Menlo Park
 - (D) STATE: California
 - (E) COUNTRY: U.S.A.
 - (F) ZIP: 94025
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: US To be assigned.
 - (B) FILING DATE: 22-SEP-1994
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Halluin, Albert P.
 - (B) REGISTRATION NUMBER: 25,227
 - (C) REFERENCE/DOCKET NUMBER: 8141-119-999
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 415-854-3660
 - (B) TELEFAX: 415-854-3694
 - (C) TELEX: 66141 PENNIE

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9629 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: unknown
 - (D) TOPOLOGY: unknown

- (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GAATTCCATC ATCAATAATA TACCTTATTT TGGATTGAAG CCAATATGAT AATGAGGGGG	60
TGGAGTTTGT GACGTGGCGC GGGGCGTGGG AACGGGGCGG GTGACGTAGT AGTGTGGCGG	120
AAGTGTGATG TTGCAAGTGT GCGGGAACAC ATGTAAGCGA CGGATGTGGC AAAAGTGACG	180

TTTTTGGTGT GCGCCGGTGT ACACAGGAAG TGACAATTTT CGCGCGGTTT TAGGCGGATG	240
TTGTAGTAAA TTTGGGCGTA ACCGAGTAAG ATTTGGCCAT TTTCGCGGGA AAACCTGAATA	300
AGAGGAAGTG AAATCTGAAT AATTTTGTGT TACTCATAGC GCGTAATATT TGTCTAGGGC	360
CGCGGGGACT TTGACCGTTT ACGTGGAGAC TCGCCCAGGT GTTTTTCTCA GGTGTTTTCC	420
CGGTTCCGGG TCAAAGTTGG CGTTTTATTA TTATAGTCTC TAGAGCTTTG CTCTTAGGAG	480
TTTCCTAATA CATCCCAAAC TCAAATATAT AAAGCATTG ACTTGTTCTA TGCCCTAGGG	540
GGCGGGGGGA AGCTAAGCCA GCTTTTTTTA ACATTTAAAA TGTTAATTCC ATTTTAAATG	600
CACAGATGTT TTTATTTTCAT AAGGGTTTCA ATGTGCATGA ATGCTGCAAT ATTCCTGTTA	660
CCAAAGCTAG TATAAATAAA AATAGATAAA CGTGGAAATT ACTTAGAGTT TCTGTCATTA	720
ACGTTTCCTT CCTCAGTTGA CAACATAAAT GCGCTGCTGA GCAAGCCAGT TTGCATCTGT	780
CAGGATCAAT TTCCCATTAT GCCAGTCATA TTAATTACTA GTCAATTAGT TGATTTTTAT	840
TTTTGACATA TACATGTGAA TGAAAGACCC CACCTGTAGG TTTGGCAAGC TAGCTTAAGT	900
AACGCCATTT TGCAAGGCAT GGAAAAATAC ATAACCTGAGA ATAGAAAAGT TCAGATCAAG	960
GTCAGGAACA GATGGAACAG CTGAATATGG GCCAAACAGG ATATCTGTGG TAAGCAGTTC	1020
CTGCCCCGGC TCAGGGCCAA GAACAGATGG AACAGCTGAA TATGGGCCAA ACAGGATATC	1080
TGTGGTAAGC AGTTCCTGCC CCGGCTCAGG GCCAAGAACA GATGGTCCCC AGATGCGGTC	1140
CAGCCCTCAG CAGTTTCTAG AGAACCATCA GATGTTTCCA GGGTGCCCCA AGGACCTGAA	1200
ATGACCCTGT GCCTTATTTG AACTAACCAA TCAGTTTCGT TCTCGCTTCT GTTCGCGCGC	1260
TTCTGCTCCC CGAGCTCAAT AAAAGAGCCC ACAACCCCTC ACTCGGGGCG CCAGTCCTCC	1320
GATTGACTGA GTCGCCCGGG TACCCGTGTA TCCAATAAAC CCTCTTGAG TTGCATCCGA	1380
CTTGTTGTTCT CGCTGTTTCT TGGGAGGGTC TCCTCTGAGT GATTGACTAC CCGTCAGCGG	1440
GGGTCTTTCA TTTGGGGGCT CGTCCGGGAT CGGGAGACCC CTGCCCAGGG ACCACCGACC	1500
CACCACCGGG AGGTAAGCTG GCCAGCAACT TATCTGTGTC TGTCCGATTG TCTAGTGTCT	1560
ATGACTGATT TTATGCGCCT GCGTCGGTAC TAGTTAGCTA ACTAGCTCTG TATCTGGCGG	1620
ACCCGTGGTG GAACTGACGA GTTCGGAACA CCCGGCCGCA ACCCTGGGAG ACGTCCCAGG	1680
GACTTCGGGG GCCGTTTTTG TGGCCCGACC TGAGTCCTAA AATCCCGATC GTTTAGGACT	1740
CTTTGGTGCA CCCCCCTTAG AGGAGGGATA TGTGGTTCTG GTAGGAGACG AGAACCTAAA	1800
ACAGTTCCCC CCTCCGTCTG AATTTTGTGT TTCGGTTTGG GACCGAAGCC GCGCCGCGCG	1860
TCTTGTCTGC TGCAGCATCG TTCTGTGTTG TCTCTGTCTG ACTGTGTTTC TGTATTTGTC	1920
TGAAAATATG GGCCCGGGCT AGACTGTTAC CACTCCCTTA AGTTTGACCT TAGGTCACTG	1980
GAAAGATGTC GAGCGGATCG CTCACAACCA GTCGGTAGAT GTCAAGAAGA GACGTTGGGT	2040
TACCTTCTGC TCTGCAGAAAT GGCCAACCTT TAACGTCGGA TGGCCGCGAG ACGGCACCTT	2100

TAACCGAGAC	CTCATCACCC	AGGTTAAGAT	CAAGGTCTTT	TCACCTGGCC	CGCATGGACA	2160
CCCAGACCAG	GTCCCCTACA	TCGTGACCTG	GGAAGCCTTG	GCTTTTGACC	CCCCTCCCTG	2220
GGTCAAGCCC	TTTGACACC	CTAAGCCTCC	GCCTCCTCTT	CCTCCATCCG	CCCCGTCTCT	2280
CCCCCTTGAA	CCTCCTCGTT	CGACCCCGCC	TCGATCCTCC	CTTTATCCAG	CCCTCACTCC	2340
TTCTCTAGGC	GCCCCCATAT	GGCCATATGA	GATCTTATAT	GGGGCACCCC	CGCCCCCTGT	2400
AAACTTCCCT	GACCCTGACA	TGACAAGAGT	TACTAACAGC	CCCTCTCTCC	AAGCTCACTT	2460
ACAGGCTCTC	TACTTAGTCC	AGCACGAAGT	CTGGAGACCT	CTGGCGGCAG	CCTACCAAGA	2520
ACAACCTGGAC	CGACCGGTGG	TACCTCACCC	TTACCGAGTC	GGCGACACAG	TGTGGGTCCG	2580
CCGACACCAG	ACTAAGAACC	TAGAACCTCG	CTGGAAAGGA	CCTTACACAG	TCCTGCTGAC	2640
CACCCCCACC	GCCCTCAAAG	TAGACGGCAT	CGCAGCTTGG	ATACACGCCG	CCCACGTGAA	2700
GGCTGCCGAC	CCCGGGGGTG	GACCATCCTC	TAGACTGCCA	TGTGGCTGCA	GAGCCTGCTG	2760
CTCTTGGGCA	CTGTGGCCTG	CAGCATCTCT	GCACCCGCC	GCTCGCCCAG	CCCCAGCAGG	2820
CAGCCCTGGG	AGCATGTGAA	TGCCATCCAG	GAGGCCCGGC	GTCTCCTGAA	CCTGAGTAGA	2880
GACACTGCTG	CTGAGATGAA	TGAAACAGTA	GAAGTCATCT	CAGAAATGTT	TGACCTCCAG	2940
GAGCCGACCT	GCCTACAGAC	CCGCCTGGAG	CTGTACAAGC	AGGGCCTGCG	GGGCAGCCTC	3000
ACCAAGCTCA	AGGGCCCCTT	GACCATGATG	GCCAGCCACT	ACAAGCAGCA	CTGCCCTCCA	3060
ACCCCGGAAA	CTTCCTGTGC	AACCCAGATT	ATCACCTTTG	AAAGTTTCAA	AGAGAACCTG	3120
AAGGACTTTC	TGCTTGTGAT	CCCCTTTGAC	TGCTGGGAGC	CAGTCCAGGA	GTGAGACCGG	3180
CCAGATGAGG	CTGGCCAAGC	CGGGGAGCTG	CTCTCTCATG	AAACAAGAGC	GGATCCTATC	3240
GCCTTCTTGA	CGAGTTCTTC	TGAGCGGGAC	TCTGGGGTTC	GAAATGACCG	ACCAAGCGAC	3300
GCCCAACCTG	CCATCACGAG	ATTTGATTC	CACCGCCGCC	TTCTATGAAA	GGTTGGGCTT	3360
CGGAATCGTT	TTCCGGGACG	CCGGCTGGAT	GATCCTCCAG	CGCGGGGATC	TCATGCTGGA	3420
GTTCTTCGCC	CACCCCAACT	TGTTTAGTCG	ACATCGATAG	ATCTGGAAGG	TGCTGAGGTA	3480
CGATGAGACC	CGCACCAGGT	GCAGACCCTG	CGAGTGTGGC	GGTAAACATA	TTAGGAACCA	3540
GCCTGTGATG	CTGGATGTGA	CCGAGGAGCT	GAGGCCCGAT	CACTTGGTGC	TGGCCTGCAC	3600
CCGCGCTGAG	TTTGGCTCTA	GCGATGAAGA	TACAGATTGA	GGTACTGAAA	TGTGTGGGCG	3660
TGGCTTAAGG	GTGGGAAAGA	ATATATAAGG	TGGGGGTCTT	ATGTAGTTTT	GTATCTGTTT	3720
TGCAGCAGCC	GCCGCGCCA	TGAGCACCAA	CTCGTTTGAT	GGAAGCATTG	TGAGCTCATA	3780
TTTGACAACG	CGCATGCCCC	CATGGGCCCG	GGTGCCTCAG	AATGTGATGG	GCTCCAGCAT	3840
TGATGGTCGC	CCCGTCTGTC	CCGCAAACTC	TACTACCTTG	ACCTACGAGA	CCGTGTCTGG	3900
AACGCCGTTG	GAGACTGCAG	CCTCCGCCGC	CGCTTCAGCC	GCTGCAGCCA	CCGCCCGCGG	3960
GATTGTGACT	GACTTTGCTT	TCCTGAGCCC	GCTTGCAAGC	AGTGCAGCTT	CCCGTTCATC	4020

CGCCCGCGAT GACAAGTTGA CGGCTCTTTT GGCACAATTG GATTCTTTGA CCCGGGAACT	4080
TAATGTCGTT TCTCAGCAGC TGTGGATCT GCGCCAGCAG GTTCTGCCC TGAAGGCTTC	4140
CTCCCCCTCCC AATGCGGTTT AAAACATAAA TAAAAAACCA GACTCTGTTT GGATTTGGAT	4200
CAAGCAAGTG TCTTGCTGTC TTTATTTAGG GGTTTTGCGC GCGCGGTAGG CCCGGGACCA	4260
GCGGTCTCGG TCGTTGAGGG TCCTGTGTAT TTTTCCAGG ACGTGGTAAA GGTGACTCTG	4320
GATGTTTCTG TACATGGGCA TAAGCCCGTC TCTGGGGTGG AGGTAGCACC ACTGCAGAGC	4380
TTCATGCTGC GGGGTGGTGT TGTAGATGAT CCAGTCGTAG CAGGAGCGCT GGGCGTGGTG	4440
CCTAAAAATG TCTTTCAGTA GCAAGCTGAT TGCCAGGGGC AGGCCCTTGG TGTAAGTGTT	4500
TACAAAGCGG TTAAGCTGGG ATGGGTGCAT ACGTGGGGAT ATGAGATGCA TCTTGACTG	4560
TATTTTTAGG TTGGCTATGT TCCCAGCCAT ATCCCTCCGG GGATTCATGT TGTGCAGAAC	4620
CACCAGCACA GTGTATCCGG TGCACTTGGG AAATTTGTCA TGTAGCTTAG AAGGAAATGC	4680
GTGGAAGAAC TTGGAGACGC CCTTGTGACC TCCAAGATTT TCCATGCATT CGTCCATAAT	4740
GATGGCAATG GGCCACGGG CGGCGGCCTG GGCGAAGATA TTTCTGGGAT CACTAACGTC	4800
ATAGTTGTGT TCCAGGATGA GATCGTCATA GGCCATTTTT ACAAAGCGCG GCGGAGGGT	4860
GCCAGACTGC GGTATAATGG TTCCATCCGG CCCAGGGGCG TAGTTACCCT CACAGATTG	4920
CATTTCCAC GCTTTGAGTT CAGATGGGGG GATCATGTCT ACCTGCGGGG CGATGAAGAA	4980
AACGGTTTCC GGGGTAGGGG AGATCAGCTG GGAAGAAAGC AGGTTCTGA GCAGCTGCGA	5040
CTTACCGCAG CCGGTGGGCC CGTAAATCAC ACCTATTACC GGGTGCAACT GGTAGTTAAG	5100
AGAGCTGCAG CTGCCGTCAT CCCTGAGCAG GGGGGCCACT TCGTTAAGCA TGTCCCTGAC	5160
TCGCATGTTT TCCCTGACCA AATCCGCCAG AAGGCGCTCG CCGCCCAGCG ATAGCAGTTC	5220
TTGCAAGGAA GCAAAGTTTT TCAACGGTTT GAGACCGTCC GCCGTAGGCA TGCTTTTGAG	5280
CGTTTGACCA AGCAGTTCCA GGCGGTCCCA CAGCTCGGTC ACCTGCTCTA CGGCATCTCG	5340
ATCCAGCATA TCTCCTCGTT TCGCGGGTTG GGGCGGCTTT CGCTGTACGG CAGTAGTCGG	5400
TGCTCGTCCA GACGGGCCAG GGTATGTCT TTCCACGGGC GCAGGGTCCT CGTCAGCGTA	5460
GTCTGGGTCA CGGTGAAGGG GTGCGCTCCG GGCTGCGCGC TGGCCAGGGT GCGCTTGAGG	5520
CTGGTCCTGC TGGTGCTGAA GCGCTGCCGG TCTTCGCCCT GCGCGTCGGC CAGGTAGCAT	5580
TTGACCATGG TGTCATAGTC CAGCCCCCTCC GCGGCGTGGC CCTTGGCGCG CAGCTTGCCC	5640
TTGGAGGAGG CGCCGCACGA GGGGCAGTGC AGACTTTTGA GGGCGTAGAG CTTGGGCGCG	5700
AGAAATACCG ATTCCGGGGA GTAGGCATCC GCGCCGCAGG CCCCAGAC GGTCTCGCAT	5760
TCCACGAGCC AGGTGAGCTC TGGCCGTTCC GGGTCAAAA CCAGGTTTCC CCCATGCTTT	5820
TTGATGCGTT TCTTACCTCT GGTTCATG AGCCGGTGTC CACGCTCGGT GACGAAAAGG	5880
CTGTCCGTGT CCCCCTATAC AGACTTGAGA GGCCTGTCCC TCGACCGATG CCCTTGAGAG	5940

CCTTCAACCC	AGTCAGCTCC	TTCCGGTGGG	CGCGGGGCAT	GACTATCGTC	GCCGCACTTA	6000
TGACTGTCTT	CTTTATCATG	CAACTCGTAG	GACAGGTGCC	GGCAGCGCTC	TGGGTCATTT	6060
TCGGCGAGGA	CCGCTTTCGC	TGGAGCGCGA	CGATGATCGG	CCTGTCGCTT	GCGGTATTCTG	6120
GAATCTTGCA	CGCCCTCGCT	CAAGCCTTCG	TCACTGGTCC	CGCCACCAA	CGTTTCGGCG	6180
AGAAGCAGGC	CATTATCGCC	GGCATGGCGG	CCGACGCGCT	GGGCTACGTC	TTGCTGGCGT	6240
TCGCGACGCG	AGGCTGGATG	GCCTTCCCCA	TTATGATTCT	TCTCGCTTCC	GGCGGCATCG	6300
GGATGCCCCG	GTTGCAGGCC	ATGCTGTCCA	GGCAGGTAGA	TGACGACCAT	CAGGGACAGC	6360
TTCAAGGATC	GCTCGCGGCT	CTTACCAGCC	TAACCTCGAT	CACTGGACCG	CTGATCGTCA	6420
CGGCGATTTA	TGCCGCCTCG	GCGAGCACAT	GGAACGGGTT	GGCATGGATT	GTAGGCGCCG	6480
CCCTATACCT	TGTCTGCCTC	CCCGCGTTGC	GTCGCGGTGC	ATGGAGCCGG	GCCACCTCGA	6540
CCTGAATGGA	AGCCGGCGGC	ACCTCGCTAA	CGGATTCACC	ACTCCAAGAA	TTGGAGCCAA	6600
TCAATTCTTG	CGGAGAACTG	TGAATGCGCA	AACCAACCCT	TGGCAGAACA	TATCCATCGC	6660
GTCCGCCATC	TCCAGCAGCC	GCACGCGGCG	CATCTCGGGC	AGCGTTGGGT	CCTGGCCACG	6720
GGTGCGCATG	ATCGTGCTCC	TGTCGTTGAG	GACCCGGCTA	GGCTGGCGGG	GTTGCCTTAC	6780
TGGTTAGCAG	AATGAATCAC	CGATACGCGA	GCGAACGTGA	AGCGACTGCT	GCTGCAAAAC	6840
GTCTGCGACC	TGAGCAACAA	CATGAATGGT	CTTCGGTTTC	CGTGTTCGT	AAAGTCTGGA	6900
AACGCGGAAG	TCAGCGCCCT	GCACCATTAT	GTTCCGGATC	TGCATCGCAG	GATGCTGCTG	6960
GCTACCCTGT	GGAACACCTA	CATCTGTATT	AACGAAGCGC	TGGCATTGAC	CCTGAGTGAT	7020
TTTTCTCTGG	TCCCCGCCGA	TCCATACCGC	CAGTTGTTTA	CCCTCACAAC	GTTCCAGTAA	7080
CCGGGCATGT	TCATCATCAG	TAACCCGTAT	CGTGAGCATC	CTCTCTCGTT	TCATCGGTAT	7140
CATTACCCCC	ATGAACAGAA	ATTCCCCCTT	ACACGGAGGC	ATCAAGTGAC	CAAACAGGAA	7200
AAAACCGCCC	TTAACATGGC	CCGCTTTATC	AGAAGCCAGA	CATTAACGCT	TCTGGAGAAA	7260
CTCAACGAGC	TGGACGCGGA	TGAACAGGCA	GACATCTGTG	AATCGCTTCA	CGACCACGCT	7320
GATGAGCTTT	ACCGCAGCTG	CCTCGCGCGT	TTCGGTGATG	ACGGTGAAAA	CCTCTGACAC	7380
ATGCAGCTCC	CGGAGACGGT	CACAGCTTGT	CTGTAAGCGG	ATGCCGGGAG	CAGACAAGCC	7440
CGTCAGGGCG	CGTCAGCGGG	TGTTGGCGGG	TGTCGGGGCG	CAGCCATGAC	CCAGTCACGT	7500
AGCGATAGCG	GAGTGTATAC	TGGCTTAACT	ATGCGGCATC	AGAGCAGATT	GTA CTGAGAG	7560
TGCACCATAT	GCGGTGTGAA	ATACCGCACA	GATGCGTAAG	GAGAAAATAC	CGCATCAGGC	7620
GCTCTTCCGC	TTCCTCGCTC	ACTGACTCGC	TGCGCTCGGT	CGTTCGGCTG	CGGCGAGCGG	7680
TATCAGCTCA	CTCAAAGGCG	GTAATACGGT	TATCCACAGA	ATCAGGGGAT	AACGCAGGAA	7740
AGAACATGTG	AGCAAAAGGC	CAGCAAAAGG	CCAGGAACCG	TAAAAAGGCC	GCGTTGCTGG	7800
CGTTTTTCCA	TAGGCTCCGC	CCCCCTGACG	AGCATCACAA	AAATCGACGC	TCAAGTCAGA	7860

GGTGGCGAAA CCCGACAGGA CTATAAAGAT ACCAGGCGTT TCCCCCTGGA AGCTCCCTCG	7920
TGCGCTCTCC TGTTCCGACC CTGCCGCTTA CCGGATACCT GTCCGCCTTT CTCCTTTCGG	7980
GAAGCGTGGC GCTTTCTCAT AGCTCACGCT GTAGGTATCT CAGTTCGGTG TAGGTTCGTTT	8040
GCTCCAAGCT GGGCTGTGTG CACGAACCCC CCGTTCAGCC CGACCGCTGC GCCTTATCCG	8100
GTAACATATCG TCTTGAGTCC AACC CGGTAA GACACGACTT ATCGCCACTG GCAGCAGCCA	8160
CTGGTAACAG GATTAGCAGA GCGAGGTATG TAGGCGGTGC TACAGAGTTC TTGAAGTGGT	8220
GGCCTAACTA CGGCTACACT AGAAGGACAG TATTTGGTAT CTGCGCTCTG CTGAAGCCAG	8280
TTACCTTCGG AAAAAGAGTT GGTAGCTCTT GATCCGGCAA ACAAACCACC GCTGGTAGCG	8340
GTGGTTTTTT TGTTTGCAAG CAGCAGATTA CGCGCAGAAA AAAAGGATCT CAAGAAGATC	8400
CTTTGATCTT TTCTACGGGG TCTGACGCTC AGTGGAACGA AAATCACGT TAAGGGATTT	8460
TGGTCATGAG ATTATCAAAA AGGATCTTCA CCTAGATCCT TTAAATTAA AAATGAAGTT	8520
TTAAATCAAT CTAAAGTATA TATGAGTAAA CTGTGCTGA CAGTTACCAA TGCTTAATCA	8580
GTGAGGCACC TATCTCAGCG ATCTGTCTAT TTCGTTTCATC CATAGTTGCC TGAATCCCCG	8640
TCGTGTAGAT AACTACGATA CGGGAGGGCT TACCATCTGG CCCCAGTGCT GCAATGATAC	8700
CGCGAGACCC ACGCTCACCG GCTCCAGATT TATCAGCAAT AAACCAGCCA GCCGGAAGGG	8760
CCGAGCGCAG AAGTGGTCTT GCAACTTTAT CCGCTCCAT CCAGTCTATT AATTGTTGCC	8820
GGGAAGCTAG AGTAAGTAGT TCGCCAGTTA ATAGTTTGCG CAACGTTGTT GCCATTGCTG	8880
CAGGCATCGT GGTGTCACGC TCGTCGTTTG GTATGGCTTC ATTCAGCTCC GGTTCCTAAC	8940
GATCAAGGCG AGTTACATGA TCCCCATGT TGTGCAAAAA AGCGGTTAGC TCCTTCGGTC	9000
CTCCGATCGT TGTCAGAAGT AAGTTGGCCG CAGTGTTATC ACTCATGGTT ATGGCAGCAC	9060
TGCATAATTC TCTTACTGTC ATGCCATCCG TAAGATGCTT TTCTGTGACT GGTGAGTACT	9120
CAACCAAGTC ATTCTGAGAA TAGTGATATG GCGCACCAGG TTGCTCTTGC CCGGCGTCAA	9180
CACGGGATAA TACCGCGCCA CATAGCAGAA CTTTAAAAGT GCTCATCATT GGAAAACGTT	9240
CTTCGGGGCG AAAACTCTCA AGGATCTTAC CGCTGTTGAG ATCCAGTTCG ATGTAACCCA	9300
CTCGTGCACC CAACTGATCT TCAGCATCTT TTACTTTTAC CAGCGTTTCT GGGTGAGCAA	9360
AAACAGGAAG GCAAAATGCC GCAAAAAGG GAATAAGGGC GACACGGAAA TGTTGAATAC	9420
TCATACTCTT CCTTTTTCAA TATTATTGAA GCATTTATCA GGGTTATTGT CTCATGAGCG	9480
GATACATATT TGAATGTATT TAGAAAAATA AACAAATAGG GGTTCCGCGC ACATTTCCCC	9540
GAAAAGTGCC ACCTGACGTC TAAGAAACCA TTATTATCAT GACATTAACC TATAAAAATA	9600
GGCGTATCAC GAGGCCCTTT CGTCTTCAA	9629

What is claimed is:

1. A chimeric adenovirus which comprises:
 - a replication deficient adenovirus genome; and
 - a DNA expression cassette comprising:
 - 5 a eucaryotic promoter and/or enhancer region;
 - nucleotide sequence corresponding to a MLV Psi-packaging site; a DNA of interest to be transcribed by said promoter; and a substantially noncoding 3' DNA which facilitates the stability, polyadenylation, or splicing
 - 10 of the transcript.
2. The chimeric adenovirus of Claim 1 wherein said DNA of interest is drawn from the group comprising:
 - granulocyte macrophage colony stimulating factor
 - 15 (GM-CSF); nerve growth factor (NGF); tyrosine hydroxylase (TH); ciliary neurotropic factor (CNTF); brain-derived neurotropic factor (BDNF); factors VIII and IX; tissue plasminogen activator (tPA); interleukins 1-2 and 4-6; tumor necrosis factor- α (TNF- α); α or γ interferons; or
 - 20 erythropoietin.
3. The chimeric adenovirus of Claim 1 wherein said DNA of interest is the gene encoding human granulocyte macrophage colony stimulating factor.
- 25 4. The chimeric adenovirus of Claim 1 wherein said DNA of interest is the gene encoding murine granulocyte macrophage colony stimulating factor.
- 30 5. A chimeric adenovirus which comprises:
 - a replication deficient adenovirus genome; and
 - a DNA expression cassette consisting essentially of an MLV LTR promoter and enhancer region; nucleotide sequence corresponding to a MLV Psi-packaging site; a
 - 35 gene encoding human granulocyte macrophage colony stimulating factor; and an SV40 polyadenylation sequence.

6. A chimeric adenovirus which comprises:
a replication deficient adenovirus genome; and
a DNA expression cassette consisting essentially of
an MLV LTR promoter and enhancer region; nucleotide
5 sequence corresponding to a MLV Psi-packaging site; a
gene encoding murine granulocyte macrophage colony
stimulating factor; and an SV40 polyadenylation sequence.
7. The use of the chimeric adenovirus of Claim 1 in the
10 treatment of mammalian disease and disorders.
8. The use of the chimeric adenovirus of Claim 2 to
transduce mammalian cells.
- 15 9. The use of the chimeric adenovirus of Claim 3 to
transduce tumor cells.
10. The use of the chimeric adenovirus of Claim 4 to
transduce tumor cells for use as anti-tumor vaccines.
20
11. A method of producing chimeric adenovirus
comprising:
the recombinatory insertion of a DNA expression
cassette into a replication deficient helper adenovirus
25 genome contained in a circular plasmid to produce a
chimeric adenovirus capable of transducing mammalian
cells.
12. The method of Claim 11 wherein said DNA expression
30 cassette comprises:
a eucaryotic promoter and/or enhancer region;
a DNA of interest to be transcribed by said
promoter; and
a 3' substantially noncoding DNA that facilitates
35 the stability, polyadenylation, or splicing of the
transcript.

13. The method of Claim 12 wherein said DNA of interest is drawn from the group comprising:

granulocyte macrophage colony stimulating factor (GM-CSF); nerve growth factor (NGF); tyrosine hydroxylase (TH); ciliary neurotropic factor (CNTF); brain-derived neurotropic factor (BDNF); factors VIII and IX; tissue plasminogen activator (tPA); interleukins 1-2 and 4-6; tumor necrosis factor- α (TNF- α); α or γ interferons; or erythropoietin.

14. The method of Claim 12 wherein said DNA of interest is the gene encoding granulocyte macrophage colony stimulating factor.

1/9




FIG.1

2/9




FIG.2

3/9

GAATTCATC ATCAATAATA TACCTTATTT TGGATTGAAG CCAATATGAT AATGAGGGGG 60
TGGAGTTTGT GACGTGGCGC GGGGCGTGGG AACGGGGCGG GTGACGTAGT AGTGTGGCGG 120
AAGTGTGATG TTGCAAGTGT GCGGGAACAC ATGTAAGCGA CGGATGTGGC AAAAGTGACG 180
TTTTTGGTGT GCGCCGGTGT ACACAGGAAG TGACAATTTT CCGCGGGTTT TAGGCGGATG 240
TTGTAGTAAA TTTGGGCGTA ACCGAGTAAG ATTTGGCCAT TTTGCGGGGA AACTGAATA 300
AGAGGAAGTG AAATCTGAAT AATTTTGTGT TACTCATAGC GCGTAATATT TGTCTAGGGC 360
CGCGGGGACT TTGACCGTTT ACGTGGAGAC TCGCCAGGT GTTTTTCTCA GGTGTTTTCC 420
GCGTTCCGGG TCAAAGTTGG CGTTTTATTA TTATAGTCTC TAGAGCTTTG CTCTTAGGAG 480
TTTCCTAATA CATCCCAAAC TCAAATATAT AAAGCATTG ACTTGTCTA TGCCCTAGGG 540
GGCGGGGGGA AGCTAAGCCA GCTTTTTTTA ACATTAAAA TGTTAATTCC ATTTTAAATG 600
CACAGATGTT TTTATTTTCA AAGGGTTTCA ATGTGCATGA ATGCTGCAAT ATTCCTGTTA 660
CCAAAGCTAG TATAAATAAA AATAGATAAA CGTGGAAATT ACTTAGAGTT TCTGTCATTA 720
ACGTTTCCTT CCTCAGTTGA CAACATAAAT GCGCTGCTGA GCAAGCCAGT TTGCATCTGT 780
CAGGATCAAT TTCCATTAT GCCAGTCATA TTAATTACTA GTCAATTAGT TGATTTTTAT 840
TTTTGACATA TACATGTGAA TGAAAGACCC CACCTGTAGG TTTGGCAAGC TAGCTTAAGT 900
AACGCCATTT TGCAAGGCAT GGAAAAATAC ATAACGTAGA ATAGAAAAGT TCAGATCAAG 960
GTCAGGAACA GATGGAACAG CTGAATATGG GCCAAACAGG ATATCTGTGG TAAGCAGTTC 1020
CTGCCCCGGC TCAGGGCCAA GAACAGATGG AACAGCTGAA TATGGGCCAA ACAGGATATC 1080
TGTGGTAAGC AGTTCCTGCC CCGGCTCAGG GCCAAGAACA GATGGTCCCC AGATGCGGTC 1140
CAGCCCTCAG CAGTTTCTAG AGAACCATCA GATGTTTCCA GGGTGCCCCA AGGACCTGAA 1200
ATGACCCTGT GCCTTATTTG AACTAACCAA TCAGTTGGCT TCTCGCTTCT GTTCGCGCGC 1260
TTCTGCTCCC CGAGCTCAAT AAAAGAGCCC ACAACCCCTC ACTCGGGGCG CCAGTCCTCC 1320
GATTGACTGA GTGCCCCGGG TACCCGTGTA TCCAATAAAC CCTCTTGACG TTGCATCCGA 1380
CTTGTGGTCT CGCTGTTCTT TGGGAGGGTC TCCTCTGAGT GATTGACTAC CCGTCAGCGG 1440
GGGTCTTTCA TTTGGGGGCT CGTCCGGGAT CGGGAGACCC CTGCCCAGGG ACCACCGACC 1500
CACCACCGGG AGGTAAGCTG GCCAGCAACT TATCTGTGTC TGTCCGATTG TCTAGTGTCT 1560
ATGACTGATT TTATGCGCCT GCGTCGGTAC TAGTTAGCTA ACTAGCTCTG TATCTGGCGG 1620
ACCCGTGGTG GAACTGACGA GTTCGGAACA CCCGGCCGA ACCCTGGGAG ACGTCCAGG 1680
GACTTCGGGG GCCGTTTTTG TGGCCCGACC TGAGTCCTAA AATCCCGATC GTTTAGGACT 1740

FIG.3A

SUBSTITUTE SHEET (RULE 26)

4/9

CTTTGGTGCA CCCCCCTAG AGGAGGGATA TGTGGTTCTG GTAGGAGACG AGAACCTAAA 1800
ACAGTTCCCG CCTCCGTCTG AATTTTGTGCT TTCGGTTTGG GACCGAAGCC GCGCCGCGCG 1860
TCTTGCTCTG TGCAGCATCG TTCTGTGTG TCTCTGTCTG ACTGTGTTTC TGTATTTGTC 1920
TGAAATATG GGCCCGGGCT AGACTGTTAC CACTCCCTTA AGTTTGACCT TAGGTCACCTG 1980
GAAAGATGTC GAGCGGATCG CTCACAACCA GTCGGTAGAT GTCAAGAAGA GACGTTGGGT 2040
TACCTTCTGC TCTGCAGAAT GGCCAACCTT TAACGTCGGA TGGCCGCGAG ACGGCACCTT 2100
TAACCGAGAC CTCATCACCC AGGTTAAGAT CAAGGTCTTT TCACCTGGCC CGCATGGACA 2160
CCCAGACCAG GTCCCTACA TCGTGACCTG GGAAGCCTTG GCTTTTGACC CCCCTCCCTG 2220
GGTCAAGCCC TTTGTACACC CTAAGCCTCC GCCTCCTCTT CCTCCATCCG CCCCCTCTCT 2280
CCCCCTTGA CCTCCTCGTT CGACCCCGCC TCGATCCTCC CTTTATCCAG CCCTCACTCC 2340
TTCTCTAGGC GCCCCCATAT GGCCATATGA GATCTTATAT GGGGCACCCC CGCCCTTGT 2400
AAACTTCCCT GACCCTGACA TGACAAGAGT TACTAACAGC CCCTCTCTCC AAGCTCACTT 2460
ACAGGCTCTC TACTTAGTCC AGCACGAAGT CTGGAGACCT CTGGCGGCAG CCTACCAAGA 2520
ACAAGTGGAC CGACCGGTGG TACCTCACCC TTACCGAGTC GGCACACAG TGTGGGTCCG 2580
CCGACACCAG ACTAAGAACC TAGAACCTCG CTGGAAAGGA CCTTACACAG TCCTGCTGAC 2640
CACCCCCACC GCCCTCAAAG TAGACGGCAT CGCAGCTTGG ATACACGCCG CCCACGTGAA 2700
GGCTGCCGAC CCGGGGGGTG GACCATCCTC TAGACTGCCA TGTGGCTGCA GAGCCTGCTG 2760
CTCTTGGGCA CTGTGGCCTG CAGCATCTCT GCACCCGCCC GCTCGCCCAG CCCCAGCAGG 2820
CAGCCCTGGG AGCATGTGAA TGCCATCCAG GAGGCCCGGC GTCTCCTGAA CCTGAGTAGA 2880
GACACTGCTG CTGAGATGAA TGAAACAGTA GAAGTCATCT CAGAAATGTT TGACCTCCAG 2940
GAGCCGACCT GCCTACAGAC CCGCCTGGAG CTGTACAAGC AGGGCCTGCG GGGCAGCCTC 3000
ACCAAGCTCA AGGGCCCCCTT GACCATGATG GCCAGCCACT ACAAGCAGCA CTGCCCTCCA 3060
ACCCCGGAAA CTTCTGTGTC AACCCAGATT ATCACCTTG AAAGTTTCAA AGAGAACCTG 3120
AAGGACTTTC TGCTTGTGAT CCCCTTTGAC TGCTGGGAGC CAGTCCAGGA GTGAGACCGG 3180
CCAGATGAGG CTGGCCAAGC CGGGGAGCTG CTCTCTCATG AAACAAGAGC GGATCCTATC 3240
GCCTTCTTGA CGAGTTCTTC TGAGCGGGAC TCTGGGGTTC GAAATGACCG ACCAAGCGAC 3300
GCCCAACCTG CCATCACGAG ATTTGATTC CACCGCCGCC TTCTATGAAA GGTGGGCTT 3360
CGGAATCGTT TTCCGGGACG CCGGCTGGAT GATCCTCCAG CGCGGGGATC TCATGCTGGA 3420
GTTCTTCGCC CACCCCAACT TGTTAGTCG ACATCGATAG ATCTGGAAGG TGCTGAGGTA 3480

FIG.3B

SUBSTITUTE SHEET (RULE 26)

5/9

CGATGAGACC CGCACCAGGT GCAGACCCTG CGAGTGTGGC GGTAACATA TTAGGAACCA 3540
GCCTGTGATG CTGGATGTGA CCGAGGAGCT GAGGCCCGAT CACTTGGTGC TGGCCTGCAC 3600
CCGCGCTGAG TTTGGCTCTA GCGATGAAGA TACAGATTGA GGTACTGAAA TGTGTGGGCG 3660
TGGCTTAAGG GTGGGAAAGA ATATATAAGG TGGGGGTCTT ATGTAGTTTT GTATCTGTTT 3720
TGCAGCAGCC GCCGCCGCCA TGAGCACCAA CTCGTTTGAT GGAAGCATTG TGAGCTCATA 3780
TTTGACAACG CGCATGCCCC CATGGGCCGG GGTGCGTCAG AATGTGATGG GCTCCAGCAT 3840
TGATGGTCGC CCCGTCTGC CCGCAAATC TACTACCTTG ACCTACGAGA CCGTGTCTGG 3900
AAGCCGTTG GAGACTGCAG CCTCCGCCG CGCTTCAGCC GCTGCAGCCA CCGCCCGCGG 3960
GATTGTGACT GACTTTGCTT TCCTGAGCCC GCTTGCAAGC AGTGCAGCTT CCCGTTTCATC 4020
CGCCCGCGAT GACAAGTTGA CGGCTCTTTT GGCACAATTG GATTCTTTGA CCCGGGAAC 4080
TAATGTCGTT TCTCAGCAGC TGTTGGATCT GCGCCAGCAG GTTCTGCCC TGAAGGCTTC 4140
CTCCCCCCC AATGCGGTTT AAAACATAAA TAAAAACCA GACTCTGTTT GGATTTGGAT 4200
CAAGCAAGTG TCTTGCTGTC TTTATTTAGG GGTTTTGCGC GCGCGGTAGG CCCGGGACCA 4260
GCGGTCTCGG TCGTTGAGGG TCCTGTGTAT TTTTCCAGG ACGTGGTAAA GGTGACTCTG 4320
GATGTTCAGA TACATGGGCA TAAGCCCGTC TCTGGGGTGG AGGTAGCACC ACTGCAGAGC 4380
TTCATGCTGC GGGGTGGTGT TGATAGTAT CCAGTCGTAG CAGGAGCGCT GGGCGTGGTG 4440
CCTAAAAATG TCTTTCAGTA GCAAGCTGAT TGCCAGGGGC AGGCCCTTGG TGTAAGTGTT 4500
TACAAAGCGG TTAAGCTGGG ATGGGTGCAT ACGTGGGGAT ATGAGATGCA TCTTGGACTG 4560
TATTTTTAGG TTGGCTATGT TCCCAGCCAT ATCCCTCCGG GGATTCATGT TGTGCAGAAC 4620
CACCAGCACA GTGTATCCGG TGCACTTGGG AAATTTGTCA TGTAGCTTAG AAGGAAATGC 4680
GTGGAAGAAC TTGGAGACGC CCTTGTGACC TCCAAGATTT TCCATGCATT CGTCCATAAT 4740
GATGGCAATG GGCCACGGG CGGCGGCCTG GGCGAAGATA TTTCTGGGAT CACTAACGTC 4800
ATAGTTGTGT TCCAGGATGA GATCGTCATA GGCCATTTTT ACAAAGCGCG GGCGGAGGGT 4860
GCCAGACTGC GGTATAATGG TTCCATCCGG CCCAGGGGCG TAGTTACCCT CACAGATTTG 3180
CATTTCCAC GCTTTGAGTT CAGATGGGGG GATCATGTCT ACCTGCGGGG CGATGAAGAA 4980
AACGGTTTCC GGGGTAGGGG AGATCAGCTG GGAAGAAAGC AGGTTCTGA GCAGCTGCCA 5040
CTTACCGCAG CCGGTGGGCC CGTAAATCAC ACCTATTACC GGGTGCAACT GGTAGTTAAG 5100
AGAGCTGCAG CTGCCGTCAT CCCTGAGCAG GGGGGCCACT TCGTTAAGCA TGTCCCTGAC 5160
TCGCATGTTT TCCCTGACCA AATCCGCCAG AAGGCGCTCG CCGCCAGCG ATAGCAGTTC 5220

FIG.3C

SUBSTITUTE SHEET (RULE 26)

6/9

TTGCAAGGAA GCAAAGTTTT TCAACGGTTT GAGACCGTCC GCCGTAGGCA TGCTTTTGAG 5280
CGTTTGACCA AGCAGTTCCA GGGCGTCCCA CAGCTCGGTC ACCTGCTCTA CGGCATCTCG 5340
ATCCAGCATA TCTCCTCGTT TCGCGGGTTG GGGCGGCTTT CGCTGTACGG CAGTAGTCGG 5400
TGCTCGTCCA GACGGGCCAG GGTCAATGTCT TTCCACGGGC GCAGGGTCCT CGTCAGCGTA 5460
GTCTGGGTCA CGGTGAAGGG GTGCGCTCCG GGCTGCGCGC TGGCCAGGGT GCGCTTGAGG 5520
CTGGTCCTGC TGGTGCTGAA GCGCTGCCGG TCTTCGCCCT GCGCGTCGGC CAGGTAGCAT 5580
TTGACCATGG TGTCATAGTC CAGCCCTCC GCGGCGTGGC CCTTGCGCGC CAGCTTGCCC 5640
TTGGAGGAGG CGCCGCACGA GGGGCAGTGC AGACTTTTGA GGGCGTAGAG CTTGGGCGCG 5700
AGAAATACCG ATTCCGGGGA GTAGGCATCC GCGCCGAGG CCCCAGAGC GGTCTCGCAT 5760
TCCACGAGCC AGGTGAGCTC TGGCCGTTCC GGGTCAAAA CCAGGTTTCC CCCATGCTTT 5820
TTGATGCGTT TCTTACCTCT GGTTCATG AGCCGGTGT CACGCTCGGT GACGAAAAGG 5880
CTGTCCGTGT CCCCCTATAC AGACTTGAGA GGCTGTCCC TCGACCGATG CCCTTGAGAG 5940
CCTTCAACCC AGTCAGCTCC TTCCGGTGGG CGCGGGGCAT GACTATCGTC GCCGCACTTA 6000
TGACTGTCTT CTTTATCATG CAACTCGTAG GACAGGTGCC GGCAGCGCTC TGGGTCATT 6060
TCGGCGAGGA CCGCTTTCCG TGGAGCGCGA CGATGATCGG CCTGTGCTT GCGGTATTCC 6120
GAATCTTGCA CGCCCTCGCT CAAGCCTTCG TCACTGGTCC CGCCACCAA CGTTTCGGCG 6180
AGAAGCAGGC CATTATCGCC GGCATGGCGG CCGACGCGCT GGGCTACGTC TTGCTGGCGT 6240
TCGGCAGCGG AGGCTGGATG GCCTTCCCA TTATGATTCT TCTCGTTCC GCGGCATCG 6300
GGATGCCCCG GTTGCAAGCC ATGCTGTCCA GGCAGGTAGA TGACGACCAT CAGGGACAGC 6360
TTCAAGGATC GCTCGCGGCT CTTACCAGCC TAACTTCGAT CACTGGACCG CTGATCGTCA 6420
CGGCGATTTA TGCCGCCTCG GCGAGCACAT GGAACGGGT GGCATGGATT GTAGCGCCG 6480
CCCTATACCT TGTCTGCCTC CCGCGTTGC GTGCGGTGC ATGGAGCCG GCCACCTCGA 6540
CCTGAATGGA AGCCGGCGG ACCTCGCTAA CGGATTCACC ACTCCAAGAA TTGGAGCCAA 6600
TCAATTCTTG CGGAGAACTG TGAATGCGCA AACCAACCCT TGGCAGAACA TATCCATCGC 6660
GTCCGCCATC TCCAGCAGCC GCACGGGCG CATCTCGGGC AGCGTTGGT CCTGGCCACG 6720
GGTGCGCATG ATCGTGCTCC TGTCGTTGAG GACCCGGCTA GGCTGGCGG GTTGCCCTAC 6780
TGGTAGCAG AATGAATCAC CGATACCGA GCGAACGTGA AGCGACTGCT GCTGCAAAAC 6840
GTCTCGGACC TGAGCAACAA CATGAATGGT CTTCCGTTT CGTGTTTCT AAAGTCTGGA 6900
AACCGGAAG TCAGCGCCCT GCACCATTAT GTTCCGGATC TGCATCGCAG GATGCTGCTG 6960

FIG.3D

SUBSTITUTE SHEET (RULE 26)

7/9

GCTACCCTGT GGAACACCTA CATCTGTATT AACGAAGCGC TGGCATTGAC CCTGAGTGAT 7020
TTTTCTCTGG TCCCGCCGCA TCCATACCGC CAGTTGTTTA CCCTCACAAC GTTCCAGTAA 7080
CCGGGCATGT TCATCATCAG TAACCCGTAT CGTGAGCATC CTCTCTCGTT TCATCGGTAT 7140
CATTACCCCC ATGAACAGAA ATTCCCCCTT ACACGGAGGC ATCAAGTGAC CAAACAGGAA 7200
AAAACCGCCC TTAACATGGC CCGCTTTATC AGAAGCCAGA CATTAAAGCT TCTGGAGAAA 7260
CTCAACGAGC TGGACGCGGA TGAACAGGCA GACATCTGTG AATCGCTTCA CGACCACGCT 7320
GATGAGCTTT ACCGCAGCTG CCTCGCGCGT TTCGGTGATG ACGGTGAAAA CCTCTGACAC 7380
ATGCAGCTCC CGGAGACGGT CACAGCTTGT CTGTAAGCGG ATGCCGGGAG CAGACAAGCC 7440
CGTCAGGGCG CGTCAGCGGG TGTGGCGGGG TGTCGGGGCG CAGCCATGAC CCAGTCACGT 7500
AGCGATAGCG GAGTGTATAC TGGCTTAAC TATGCGGCATC AGAGCAGATT GACTGAGAG 7560
TGCACCATAT GCGGTGTGAA ATACCGCACA GATGCGTAAG GAGAAAATAC CGCATCAGGC 7620
GCTCTTCCGC TTCCTCGCTC ACTGACTCGC TCGGCTCGGT CGTTCGGCTG CGGCGAGCGG 7680
TATCAGCTCA CTCAAAGGCG GTAATACGGT TATCCACAGA ATCAGGGGAT AACGCAGGAA 7740
AGAACATGTG AGCAAAAGGC CAGCAAAAGG CCAGGAACCG TAAAAAGGCC GCGTTGCTGG 7800
CGTTTTTCCA TAGGCTCCGC CCCCCTGACG AGCATCACAA AAATCGACGC TCAAGTCAGA 7860
GGTGGCGAAA CCCGACAGGA CTATAAGAT ACCAGGCGTT TCCCCTGGA AGCTCCCTCG 7920
TGCGCTCTCC TGTTCGACC CTGCCGTTA CCGGATACCT GTCCGCCTTT CTCCCTTCGG 7980
GAAGCGTGGC GCTTTCTCAT AGCTCACGCT GTAGGTATCT CAGTTCGGTG TAGGTCGTTT 8040
GCTCCAAGCT GGGCTGTGTG CACGAACCCC CCGTTCAGCC CGACCGCTGC GCCTTATCCG 8100
GTAACATCG TCTTGAGTCC AACCCTGTAA GACACGACTT ATCGCCACTG GCAGCAGCCA 8160
CTGGTAACAG GATTAGCAGA GCGAGGTATG TAGGCGGTGC TACAGAGTTC TTGAAGTGGT 8220
GGCCTAACTA CGGCTACACT AGAAGGACAG TATTTGGTAT CTGCGCTCTG CTGAAGCCAG 8280
TTACCTTCGG AAAAAGAGTT GGTAGCTCTT GATCCGGCAA ACAAACCACC GCTGGTAGCG 8340
GTGGTTTTTT TGTTCGAAG CAGCAGATTA CCGCGAGAAA AAAAGGATCT CAAGAAGATC 8400
CTTTGATCTT TTCTACGGGG TCTGACGCTC AGTGGAAACGA AAATCACGT TAAGGGATTT 8460
TGGTCATGAG ATTATCAAAA AGGATCTTCA CCTAGATCCT TTAAATTAA AAATGAAGTT 8520
TTAAATCAAT CTAAGTATA TATGAGTAA CTTGGTCTGA CAGTTACCAA TGCTTAATCA 8580
GTGAGGCACC TATCTCAGCG ATCTGTCTAT TTCGTTTCATC CATAGTTGCC TGAATCCCG 8640
TCGTGTAGAT AACTACGATA CGGGAGGGCT TACCATCTGG CCCCAGTGCT GCAATGATAC 8700

FIG.3E

SUBSTITUTE SHEET (RULE 26)

8/9

CGCGAGACCC ACGCTACCG GCTCCAGATT TATCAGCAAT AAACCAGCCA GCCGGAAGGG 8760
CCGAGCGCAG AAGTGGTCCT GCAACTTTAT CCGCTCCAT CCAGTCTATT AATTGTTGCC 8820
GGGAAGCTAG AGTAAGTAGT TCGCCAGTTA ATAGTTTGCG CAACGTTGTT GCCATTGCTG 8880
CAGGCATCGT GGTGTCACGC TCGTCGTTG GTATGGCTTC ATTCAGCTCC GGTTCCTAAC 8940
GATCAAGGCG AGTTACATGA TCCCCATGT TGTGCAAAAA AGCGGTAGC TCCTTCGGTC 9000
CTCCGATCGT TGTGAGAAGT AAGTTGGCCG CAGTGTTATC ACTCATGGTT ATGGCAGCAC 9060
TGCATAATTC TCTTACTGTC ATGCCATCCG TAAGATGCTT TTCTGTGACT GGTGAGTACT 9120
CAACCAAGTC ATTCTGAGAA TAGTGTATGC GCGGACCGAG TTGCTCTTGC CCGGCGTCAA 9180
CACGGGATAA TACCGCGCCA CATAGCAGAA CTTTAAAAGT GTCATCATT GGAAAACGTT 9240
CTTCGGGGCG AAAACTCTCA AGGATCTTAC CGCTGTTGAG ATCCAGTTCG ATGTAACCCA 9300
CTCGTGCACC CAACTGATCT TCAGCATCTT TTACTTTCAC CAGCGTTTCT GGGTGAGCAA 9360
AAACAGGAAG GCAAAATGCC GCAAAAAAGG GAATAAGGGC GACACGGAAA TGTGAATAC 9420
TCATACTCTT CCTTTTTCAA TATTATTGAA GCATTATCA GGGTTATTGT CTCATGAGCG 9480
GATACATATT TGAATGTATT TAGAAAAATA AACAAATAGG GGTCCGCGC ACATTTCCCC 9540
GAAAAGTGCC ACCTGACGTC TAAGAAACCA TTATTATCAT GACATTAACC TATAAAAATA 9600
GGCGTATCAC GAGGCCCTTT CGTCTTCAA 9629

FIG.3F

SUBSTITUTE SHEET (RULE 26)




FIG. 4A




FIG. 4B




FIG. 5





FIG. 6

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C12N 15/86, C07K 14/535	A3	(11) International Publication Number: WO 96/09399 (43) International Publication Date: 28 March 1996 (28.03.96)
(21) International Application Number: PCT/US95/11537 (22) International Filing Date: 12 September 1995 (12.09.95) (30) Priority Data: 311,485 23 September 1994 (23.09.94) US (71) Applicant: SOMATIX THERAPY CORPORATION [US/US]; Suite 100, 950 Marina Village Parkway, Alameda, CA 94501 (US). (72) Inventors: SHANKARA, Srinivas; Apartment E, 2255 San Jose Avenue, Alameda, CA 94501 (US). DWARKI, Varavani; Apartment N, 1175 Broadway Street, Alameda, CA 94501 (US). NUJAR, Tarlochan; 946 Foxfire Drive, Maneca, CA 95336 (US). (74) Agents: HALLUIN, Albert, P. et al.; Pennie & Edmonds, 1155 Avenue of the Americas, New York, NY 10036 (US).	(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i> (88) Date of publication of the international search report: 18 July 1996 (18.07.96)	

(54) Title: CHIMERIC ADENOVIRUS FOR GENE DELIVERY



(57) Abstract

Chimeric adenovirus capable of transducing mammalian cells with DNA of interest are disclosed. The chimeric adenovirus are useful for the delivery of cloned genes into an individual and are therefore also useful for treating mammalian genetic diseases and disorders.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

INTERNATIONAL SEARCH REPORT

Inter: nal Application No
PCT/US 95/11537A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/86 C07K14/535

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO,A,93 03163 (FONDATION NATIONALE DE TRANSFUSION SANGUINE) 18 February 1993 see page 4, line 19 - page 8, line 18; example 5	1-14
A	--- EUROPEAN JOURNAL OF NEUROSCIENCE, vol. 5 , no. 10 , 1 October 1993, pages 1287-1291, XP002002600 C.CAILLAUD ET AL.: "Adenoviral vector as a gene delivery system into cultured rat neuronal and glial cells" --- -/--	1-14

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *A* document member of the same patent family

Date of the actual completion of the international search

9 May 1996

Date of mailing of the international search report

23.05.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Cupido, M

INTERNATIONAL SEARCH REPORT

Inter. Application No.
PCT/US 95/11537

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 91, 12 April 1994, WASHINGTON US, pages 3054-3057, XP002002601 S-H CHEN ET AL.: "Gene therapy for brain tumors:Regression of experimental gliomas by adenovirus-mediated gene transfer in vivo." see paragraph bridging left and right columns on page 3054 -----</p>	1-14

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/11537

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 7, 10
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 7 and 10 are directed to a method of treatment of the human body the search has been carried out and based on the alleged effects of the composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

information on patent family members

PCT/US 95/11537

18-05-94
13-10-94